

Table 2.2

Age-Adjusted SEER Cancer Incidence Rates^a
 By International Classification of Childhood Cancer (ICCC)^b Selected Group and Subgroup^c and Year of Diagnosis
 Excluding benign brain and myelodysplastic syndromes
 All Races, Males and Females, Ages 0-19

ICCC Group and Subgroup	1975-2018	1975-1986	1987-1996	1997-2008	2009-2018
All ICCC Groups Combined	165.8	146.4	159.3	168.1	189.4
I Leukemia	40.8	35.5	38.2	42.8	46.3
I(a) Lymphoid leukemia	29.9	26.1	28.6	31.4	33.2
I(b) Acute myeloid leukemia	7.4	6.2	6.9	8.0	8.2
II Lymphomas and reticuloendothelial neoplasms	26.3	24.6	24.4	24.2	32.2
II(a) Hodgkin lymphoma	13.0	14.3	13.5	11.8	12.7
II(b,c,e) Non-Hodgkin lymphoma					
III CNS & misc intracranial & intraspinal neoplasms	28.9	24.1	29.9	29.7	31.8
III(a) Ependymomas and choroid plexus tumor	2.5	1.9	2.6	2.5	3.1
III(b) Astrocytoma	14.6	12.1	15.7	14.7	16.2
III(c) Intracranial and intraspinal embryonal tumors	5.6	4.5	5.7	6.2	6.1
III(d) Other gliomas	5.3	4.6	5.1	5.6	5.7
IV Neuroblastoma and other peripheral nervous cell tumors	8.6	8.2	8.5	8.7	8.9
IV(a) Neuroblastoma and ganglioneuroblastoma	8.3	8.0	8.2	8.5	8.6
V Retinoblastoma	3.1	2.8	3.2	3.2	3.1
VI Renal tumors	6.7	6.5	6.8	6.7	6.7
VI(a) Nephroblastoma and other nonepithelial renal tumor	6.1	6.2	6.4	6.0	6.0
VII Hepatic tumors	2.2	1.4	1.8	2.5	2.9
VII(a) Hepatoblastoma	1.6	0.9	1.4	1.8	2.2
VIII Malignant bone tumors	8.7	8.4	8.8	8.6	9.2
VIII(a) Osteosarcoma	4.8	4.3	4.9	4.7	5.4
VIII(c) Ewing tumor and related sarcomas of bone	3.0	3.2	2.8	2.9	3.0
IX Soft tissue and other extraosseous sarcomas	11.6	10.7	10.9	12.4	12.1
IX(a) Rhabdomyosarcoma	4.4	4.2	4.7	4.6	4.1
X Germ cell & trophoblastic tumors & neoplasms of gonads	10.7	9.4	10.8	11.0	11.5
X(a) Intracranial and intraspinal germ-cell tumor	1.5	1.0	1.6	1.7	1.9
X(c) Malignant gonadal germ cell tumor	6.8	6.0	6.7	7.2	7.1
XI Other malignant epithelial neoplasms and melanomas ^d	17.6	14.0	15.2	17.4	23.8
XI(b) Thyroid carcinoma	6.9	5.0	5.2	6.4	11.0
XI(d) Malignant melanoma	5.0	4.2	5.3	6.0	4.7

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Rates are per 1,000,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b International Classification of Childhood Cancer is based on ICD-O-3. Steliarova-Foucher E, Colombet M, Ries LAG, Moreno F, Dolya A, Shin HY, Hesselting P, Stiller CA. International Incidence of Childhood Cancer, Volume III. Lyon: International Agency for Research on Cancer, In press.

^c Classifications are shown for invasive cases only except as noted. For ICCC groups and subgroups that include *in situ* behavior, only invasive cases are shown except as noted.

^d Includes *in situ* urinary bladder tumors.

- Statistic could not be calculated. Rate based on less than 16 cases for the time interval.